




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 401/06, 401/14, 409/14, A61K 31/47		A1	(11) International Publication Number: WO 97/49695
			(43) International Publication Date: 31 December 1997 (31.12.97)
(21) International Application Number: PCT/EP97/03160		(81) Designated States: JP, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(22) International Filing Date: 17 June 1997 (17.06.97)		Published With international search report.	
(30) Priority Data: 9613261.8 25 June 1996 (25.06.96) GB 9613262.6 25 June 1996 (25.06.96) GB			
(71) Applicant (for all designated States except US): SMITHKLINE BEECHAM P.L.C. [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB).			
(72) Inventors; and			
(75) Inventors/Applicants (for US only): FORBES, Ian, Thomson [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). RAHMAN, Shirley, Katherine [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB).			
(74) Agent: WATERS, David, Martin; SmithKline Beecham, Corporate Intellectual Property, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).			

(54) Title: SULFONAMIDE DERIVATIVES AS 5HT7 RECEPTOR ANTAGONISTS



(57) Abstract

The invention relates to sulphonamide compounds of formula (I) or a salt thereof having pharmacological activity, processes for their preparation, to compositions containing them and to their use in the treatment of CNS disorders, wherein: Ar is an optionally substituted mono- or bicyclic aromatic or heteroaromatic ring; Ar' is an optionally substituted mono- or bicyclic aromatic or heteroaromatic ring; R¹ is C₁₋₆alkyl or together with the group R³ form a 5-8 membered ring containing one or two heteroatoms optionally substituted by C₁₋₆alkyl; R² is hydrogen or C₁₋₆alkyl; R³ is hydrogen, C₁₋₆alkyl or together with the group R¹ form a 5-8 membered ring containing one or two heteroatoms optionally substituted by C₁₋₆alkyl; R⁴ is hydrogen or C₁₋₆alkyl; R⁵ and R⁶ are independently hydrogen or C₁₋₆alkyl; p is 1, 2 or 3; q is 1 to 3; and r is 1 or 2.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.


AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

SULFONAMIDE DERIVATIVES AS 5HT₇ RECEPTOR ANTAGONISTS

This invention relates to compounds having pharmacological activity, processes for their preparation, to compositions containing them and to their use in the treatment of CNS disorders.

EPA 0 021 580 and EPA 0 076 072 describe sulphonamide derivatives which are disclosed as having antiarrhythmic activity. A structurally distinct class of compounds has now been discovered, which have been found to have 5HT₇ receptor antagonist activity. 5HT₇ receptor antagonists are believed to be of potential use in the treatment of certain CNS disorders such as anxiety, depression, sleep disorders, and schizophrenia.

The present invention therefore provides, in a first aspect, a compound of formula (I) or a salt thereof:



(I)

wherein:

- Ar is an optionally substituted mono- or bicyclic aromatic or heteroaromatic ring;
- Ar' is an optionally substituted mono- or bicyclic aromatic or heteroaromatic ring;
- R¹ is C₁₋₆alkyl or together with the group R³ form a 5-8 membered ring containing one or two heteroatoms optionally substituted by C₁₋₆alkyl;
- R² is hydrogen or C₁₋₆alkyl;
- R³ is hydrogen, C₁₋₆alkyl or together with the group R¹ form a 5-8 membered ring containing one or two heteroatoms optionally substituted by C₁₋₆alkyl;
- R⁴ is hydrogen or C₁₋₆alkyl;
- R⁵ and R⁶ are independently hydrogen or C₁₋₆alkyl;
- p is 1, 2 or 3;
- q is 1, 2 or 3; and

r is 1 or 2.

C₁₋₆ Alkyl groups, whether alone or as part of another group, may be straight chain or branched.

Optional substituents for aromatic and heteroaromatic groups include

5 C₁₋₆ alkyl optionally substituted by NR⁷R⁸, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ alkylthio, cyano, nitro, halogen, CF₃, C₂F₅, NR⁷R⁸, CONR⁷R⁸, NR⁷COR⁸, S(O)_pNR⁷R⁸, CHO, OCF₃, SCF₃, SOR⁹, SO₂R⁹, OSO₂R⁹, COR⁹, CH₂OR⁹, CO₂R⁹ or OR⁹ where p is 1 or 2 and R⁷, R⁸ and R⁹ are independently hydrogen, C₁₋₆ alkyl, optionally substituted aryl or optionally substituted arylC₁₋₆alkyl. More than

10 one substituent can be present and in the case of multiple substituents these can be the same or different.

Preferably Ar is an optionally substituted bicyclic aromatic group. Most preferably Ar is naphthyl.

Preferably Ar' is an optionally substituted monocyclic aromatic group. Most

15 preferably Ar' is phenyl.

Examples of groups where R¹ is C₁₋₆alkyl are ethyl and most preferably methyl. When groups R¹ and R³ are combined to form a heterocycle the preferred examples have 5 or most preferably 6 membered rings.

Preferably R² is hydrogen or methyl;

20 The preferred group when R³ is C₁₋₆alkyl is methyl. When groups R¹ and R³ are combined to form a heterocycle the preferred examples have 5 or most preferably 6 membered rings.

Preferably R⁴ is hydrogen;

Preferably R⁵ and R⁶ are hydrogen;

25 Preferably q and r have values such that they form part of a 5- or 6-membered ring. Most preferably q and r have values such that they form part of a 6-membered ring i.e. the sum of q and r is 3.

Particular compounds of the invention include:

2-(2-[1-Naphthalene-1-sulfonyl]piperidin-2-yl)-ethyl-1,2,3,4-tetrahydroisoquinoline

30 2-(2-(3-Chloro-4-methylphenyl)-piperidin-2-yl)-ethyl-1,2,3,4 tetrahydroisoquinoline and pharmaceutically acceptable salts thereof.

The compounds of the formula (I) can form acid addition salts with acids, such as conventional pharmaceutically acceptable acids, for example maleic, hydrochloric, hydrobromic, phosphoric, acetic, fumaric, salicylic, citric, lactic, mandelic, tartaric

35 and methanesulfonic.

Compounds of formula (I) may also form solvates such as hydrates, and the invention also extends to these forms. When referred to herein, it is understood that the term 'compound of formula (I)' also includes these forms.

Certain compounds of formula (I) are capable of existing in stereoisomeric forms including diastereomers and enantiomers and the invention extends to each of these stereoisomeric forms and to mixtures thereof including racemates. The different stereoisomeric forms may be separated one from the other by the usual methods, or any given isomer may be obtained by stereospecific or asymmetric synthesis. The invention also extends to any tautomeric forms and mixtures thereof.


The present invention also provides a process for the preparation of a compound of formula (I) or a pharmaceutically acceptable salt thereof, which process comprises

(a) the coupling of a compound of formula (II):




(II)

in which Ar is as defined in formula (I) and L is a leaving group with a compound of formula (III):




(III)

in which p, q, r, R², R³, R⁴, R⁵, R⁶ and Ar' are as defined in formula (I);
or (b) the coupling of a compound of formula (IV):



(IV)

in which Ar, p, R¹, R², R³, R⁵ and R⁶ are as defined in formula (I) and L¹ is a leaving group with a compound of formula (V):



in which q, r, R⁴ and Ar' are as defined in formula (I) and optionally thereafter (a) or (b):

- forming a pharmaceutically acceptable salt.

Suitable leaving groups L and L¹ include halogen, in particular chloro. The reaction of a compounds of formulae (II) and (III) is preferably carried out in an inert solvent such as dichloromethane optionally in the presence of a base such as triethylamine.

Compounds of formulae (II), (III), (IV) and (V) are either commercially available or may be prepared according to known methods or analogous to known methods. Novel compounds of formulae (II), (III), (IV) and (V) form a further aspect of the invention.

Pharmaceutically acceptable salts may be prepared conventionally by reaction with the appropriate acid or acid derivative.

Compounds of formula (I) and their pharmaceutically acceptable salts have 5HT₇ receptor antagonist activity and are believed to be of potential use for the treatment or prophylaxis of CNS disorders such as anxiety, depression, sleep disorders, including instances of Circadian rhythm and schizophrenia.

Thus the invention also provides a compound of formula (I) or a pharmaceutically acceptable salt thereof, for use as a therapeutic substance, in particular in the treatment or prophylaxis of the above disorders.

The invention further provides a method of treatment or prophylaxis of the above disorders, in mammals including humans, which comprises administering to the sufferer a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof.

In another aspect, the invention provides the use of a compound of formula (I) or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for the treatment or prophylaxis of the above disorders.

The present invention also provides a pharmaceutical composition, which comprises a compound of formula (I) or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

5 A pharmaceutical composition of the invention, which may be prepared by admixture, suitably at ambient temperature and atmospheric pressure, is usually adapted for oral, parenteral or rectal administration and, as such, may be in the form of tablets, capsules, oral liquid preparations, powders, granules, lozenges, reconstitutable powders, injectable or infusable solutions or suspensions or suppositories. Orally administrable compositions are generally preferred.

10 Tablets and capsules for oral administration may be in unit dose form, and may contain conventional excipients, such as binding agents, fillers, tableting lubricants, disintegrants and acceptable wetting agents. The tablets may be coated according to methods well known in normal pharmaceutical practice.

15 Oral liquid preparations may be in the form of, for example, aqueous or oily suspension, solutions, emulsions, syrups or elixirs, or may be in the form of a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, emulsifying agents, non-aqueous vehicles (which may include edible oils), preservatives, and, if desired, conventional flavourings or colourants.

20 For parenteral administration, fluid unit dosage forms are prepared utilising a compound of the invention or pharmaceutically acceptable salt thereof and a sterile vehicle. The compound, depending on the vehicle and concentration used, can be either suspended or dissolved in the vehicle. In preparing solutions, the compound can be dissolved for injection and filter sterilised before filling into a suitable vial or
25 ampoule and sealing. Advantageously, adjuvants such as a local anaesthetic, preservatives and buffering agents are dissolved in the vehicle. To enhance the stability, the composition can be frozen after filling into the vial and the water removed under vacuum. Parenteral suspensions are prepared in substantially the same manner, except that the compound is suspended in the vehicle instead of being
30 dissolved, and sterilization cannot be accomplished by filtration. The compound can be sterilised by exposure to ethylene oxide before suspension in a sterile vehicle. Advantageously, a surfactant or wetting agent is included in the composition to facilitate uniform distribution of the compound.

35 The composition may contain from 0.1% to 99% by weight, preferably from 10 to 60% by weight, of the active material, depending on the method of administration.

The dose of the compound used in the treatment of the aforementioned disorders will vary in the usual way with the seriousness of the disorders, the weight

of the sufferer, and other similar factors. However, as a general guide suitable unit doses may be 0.05 to 1000 mg, more suitably 0.05 to 20.0 mg, for example 0.2 to 5 mg; and such unit doses may be administered more than once a day, for example two or three a day, so that the total daily dosage is in the range of about 0.5 to 100 mg; and such therapy may extend for a number of weeks or months.

When administered in accordance with the invention, no unacceptable toxicological effects are expected with the compounds of the invention.

The following Descriptions and Examples illustrate the preparation of compounds of the invention.

Description 1

2-(2-Chloroethyl)-1-(naphthalene-1-sulfonyl)piperidine (D1)

To a solution of 1-naphthalene sulfonyl chloride (26.64g) in toluene (300 ml) was added 2-piperidine ethanol (8.99g) and diisopropylethylamine (26.8 ml). The mixture was heated to reflux overnight. After cooling to room temperature the solvent was concentrated *in vacuo* and the residue chromatographed on silica eluting with 50% ethyl acetate and petroleum ether (bp 60-80). The title compound was isolated as an oil, which solidified on standing (12.5g, 53%). MH^+ 338.

2-[1-(naphthalene-1-sulfonyl)-piperidin-2-yl]ethanol the more polar product was isolated as an oil (9.8g, 44%).

Description 2

2-(2-Hydroxyethyl)-piperidine-1-carboxylic acid benzyl ester (D2)

2-Piperidine ethanol (39.6 ml, 0.31 mol) was dissolved in 5M aq NaOH (62 ml, 0.31 mol) and dioxan (100 ml). The mixture was cooled to 0°C and treated with benzyl chloroformate (45.3 ml, 0.32 mol) and 5M NaOH (62 ml, 0.31 mol). Stirring was continued at room temp. for 18 hours. Dioxan was removed *in vacuo* and the aqueous phase extracted with ether before acidifying with 5N HCl and extracting again with ether. The organic phase was dried and evaporated *in vacuo* and the residue purified by chromatography on silica gel to give the title compound (77.2g, 95%). MH^+ 264.

Description 3

2-[2-(3,4-Dihydro-1H-isoquinolin-2-yl)ethyl]piperidine-1-carboxylic acid benzyl ester (D3)

To a solution of 2-(2-hydroxyethyl)-piperidine-1-carboxylic acid benzyl ester (D2, 20g, 76 mmol) in dichloromethane (400 ml) was added triethylamine (21 ml, 152 mmol) and methanesulfonic anhydride (20g, 114 mmol) at 0°C. Stirring was continued for 1 hr. The reaction mixture was washed with sat. aq. NaHCO₃, dried and concentrated to afford the crude mesylate. This was dissolved in dichloromethane (150 ml) and treated with 1,2,3,4-tetrahydroisoquinoline (21 ml, 167 mmol). Stirring was continued for 48 hrs. The reaction mixture was partitioned between H₂O and CH₂Cl₂. The organic phase was dried and concentrated and the residue purified by chromatography on silica gel to afford the title compound (13g, 49%). MH⁺ 379.

Description 4

2-(2-Piperidin-2-yl ethyl)-1,2,3,4-tetrahydroisoquinoline (D4)


The protected amine 2-[2-(3,4-dihydro-1H-isoquinolin-2-yl)ethyl]piperidine-1-carboxylic acid benzyl ester (D3, 14g, 0.037 mol) was dissolved in ethanol (300 ml) and hydrogenated over 10% Pd/C for 18 hours to afford the title compound (9g, 100%). MH⁺ 245.

Example 1

2-(2-[1-Naphthalene-1-sulfonyl]piperidin-2-yl)-ethyl)-1,2,3,4-tetrahydroisoquinoline (E1)

To a solution of 2-(2-chloroethyl)-1-(naphthalene-1-sulfonyl)-piperidine (D1, 250 mg) in acetonitrile (20 ml) was added sodium iodide (12 mg), potassium carbonate (108 mg) and 1,2,3,4-tetrahydroisoquinoline (117 µl). The mixture was heated at reflux overnight. After cooling to room temperature the solvent was removed under reduced pressure. The residue was chromatographed on silica eluting with 5% methanol in dichloromethane to afford pure title compound as a foam (127 mg, 40%) MH⁺ 435.

Examples E2-12 were prepared using the procedure outlined in Example 1 using 2-(2-chloroethyl)-1-(naphthalene-1-sulfonyl)piperidine (D1) and an appropriately substituted 1,2,3,4 tetrahydroisoquinoline



Example	R	MH ⁺
2	7-Nitro	480
3	7-Methylsulfonyl	513
4	7-Sulphamoyl	514
5	6-Methoxy	465
6	7-Methoxy	465
7	5-Acetylamino	492
8	8-Chloro	469/471
9	6,7-Dimethoxy	495
10	7-Benzoyl	539
11	7-Phenoxy	527
12	8-Phenyl	511

Example 13

5 (+/-) 7-Amino-2-(2-[1-(naphthalene-1-sulfonyl)piperidin-2-yl]ethyl)-1,2,3,4-tetrahydroisoquinoline (E13)

10 A warm solution of 7-nitro-2-(2-[1-(naphthalene-1-sulfonyl)piperidin-2-yl]ethyl)-1,2,3,4-tetrahydroisoquinoline (E2) (22 mmol) in methanol (300 ml) was added to a solution of ammonium chloride (6.0g) in water (100 ml) containing iron powder (3.8g). The mixture was heated to reflux for 5 hours. The mixture was filtered whilst hot and the methanol was subsequently removed *in vacuo*. The crude product was crystallised from ethyl acetate/methanol to give the title compound (5.9g, 58%) mpt 212-218°C. MH⁺ 450.


15 **Example 14**

(+/-) 7-Bromo-2-(2-[1-(naphthalene-1-sulfonyl)piperidin-2-yl]ethyl)-1,2,3,4-tetrahydroisoquinoline (E14)

To a solution of 7-amino-2-(2-[1-(naphthalene-1-sulfonyl)piperidin-2-yl]ethyl)-1,2,3,4-tetrahydroisoquinoline (E13) (4g, 8.9 mmol) in 45% HBr at 0°C was added sodium nitrite (0.65g) in water (5 ml). The resulting mixture was added portionwise to a solution of copper (I) bromide (0.75g) in hydrobromic acid (3 ml). When effervescence had ceased, the mixture was partitioned between 40% aqueous NaOH and CH₂Cl₂/MeOH. The organic phase was separated, filtered, dried and evaporated. The residue was purified on silica to afford the title compound (2.73g, 60%). MH⁺ 513/515.

10 **Examples E15-24 were prepared by the following generic procedure:**

A mixture of (+/-) 7-bromo-2-(2-[1-naphthalene-1-sulfonyl)piperidin-2-yl]ethyl)-1,2,3,4-tetrahydroisoquinoline (E14) (1 mmol), and an arylboronic acid (1 mmol), sodium carbonate (4 mmol), tetrakis(triphenylphosphine)palladium (0) (0.03 mmol) in 1,2 DME and water (1:1, 20 ml) was heated to reflux for 3 hrs. After removal of the DME, the product was extracted with CH₂Cl₂. The organic phase was washed with brine, dried and concentrated. The crude product was converted to its hydrochloride salt with 1M HCl in Et₂O.



Example	Ar	MH ⁺
15	Phenyl	511
16	2-Methylphenyl	525
17	4-Methylphenyl	525
18	4-Trifluoromethylphenyl	579
19	3-Trifluoromethylphenyl	579
20	3-Methoxyphenyl	541
21	3-Pyridyl	512
22	2-Methoxyphenyl	541
23	4-Methoxyphenyl	541
24	3-Chlorophenyl	545/547

Example 25

(+/-) 2-(2-[1-(Naphthalene-1-sulfonyl)piperidin-2-yl]ethyl)-1,2,3,4-tetrahydro-7-trifluoromethyl isoquinoline (E25)

5

A mixture of 7-bromo-2-(2-[1-naphthalene-1-sulfonyl)piperidin-2-yl]ethyl-1,2,3,4-tetrahydroisoquinoline (E14) (0.5g, 0.97 mmol), potassium trifluoroacetate (3 mmol) and copper (I) iodide (0.56g) in toluene (3 ml) and DMF (5 ml) was heated to 135°C, for 24 hrs. The mixture was filtered and evaporated and the residue partitioned
10 between ethyl acetate and water. The organic phase was washed with brine, dried and concentrated *in vacuo*. Purification by chromatography on silica gave the title compound. MH^+ 503.

Example 26

15 (+/-) 6-Hydroxy-2-(2-[1-(naphthalene-1-sulfonyl)piperidin-2-yl]ethyl)-1,2,3,4-tetrahydroisoquinoline (E26)

Boron tribromide (1M in CH_2Cl_2) (6.6 ml) was added to a solution of 6-methoxy-2-(2-[1-(naphthalene-1-sulfonyl)piperidin-2-yl]ethyl)-1,2,3,4-tetrahydroisoquinoline (E5)
20 (0.5g, 1.1 mmol) in dichloromethane (20 ml). The mixture was stirred at room temp. overnight and then poured onto ice cold conc. ammonia solution. The organic phase was washed with brine, dried and concentrated. Purification by chromatography on silica gel afforded the title compound (0.2g, 40%). MH^+ 451.

Example 27

25 (+/-) 6-Phenyl-2-(2-[1-(naphthalene-1-sulfonyl)piperidin-2-yl]ethyl)-1,2,3,4-tetrahydroisoquinoline (E27)

To a solution of (+/-) 6-hydroxy-2-(2-[1-(naphthalene-1-sulfonyl)piperidin-2-yl]ethyl)-1,2,3,4-tetrahydroisoquinoline (E26) (0.16g, 0.35 mmol) in pyridine (2 ml)
30 at 0°C was added trifluoromethane sulfonic anhydride (0.39 mmol). After stirring for 48 hrs the mixture was concentrated, dissolved in dichloromethane, washed with water, dried and concentrated to give the 6-trifluoromethane sulphonate derivative (0.18g, 66%). This compound was converted to the title compound using
35 phenylboronic acid and the method described in Example 15. MH^+ 511.

Example 28

(+/-) 5-Amino-2-(2-[1-(naphthalene-1-sulfonyl)piperidin-2-yl]ethyl)-1,2,3,4-tetrahydroisoquinoline (E28)

5 A mixture of the 5-acetylamino compound (Example 7) (1.1g, 2.2 mmol) and 20% aqueous NaOH (3.5 ml) in EtOH (20 ml) was heated to reflux for 48 hrs. The mixture was concentrated and the residue dissolved in CH₂Cl₂. The solution was extracted twice with 5M HCl. The acidic extracts were basified with 40% NaOH and extracted with CH₂Cl₂. The organic extract was concentrated and the residue purified by chromatography on silica gel to afford the title compound (0.51g, 51%). MH⁺ 450.

10

Example 29

(+/-) 5-Bromo-2-(2-[1-(naphthalene-1-sulfonyl)piperidin-2-yl]ethyl)-1,2,3,4-tetrahydroisoquinoline (E29)

15 The 5-amino compound (Example 28) (0.43g, 0.96 mmol) was converted to the title compound using the method described in Example 14 (0.31g, 63%). MH⁺ 513/515.

Example 30

20 **(+/-) 2-(2-[1-(Naphthalene-1-sulfonyl)piperidine-2-yl]ethyl)-5-phenyl-1,2,3,4-tetrahydroisoquinoline (E30)**

The title compound was prepared according to the procedure outlined in Example 15 using (+/-) 5-bromo-2-(2-[1-(naphthalene-1-sulfonyl)piperidin-2-yl]ethyl)-1,2,3,4-tetrahydroisoquinoline (E29) and phenyl boronic acid. MH⁺ 511.

25

Example 31

(+/-) 7-Benzyloxy-2-(2-[1-(naphthalene-1-sulfonyl)piperidin-2-yl]ethyl)-1,2,3,4-tetrahydroisoquinoline (E31)


30 The 7-methoxy compound (Example 6) (0.93g, 2 mmol) was treated with BBr₃ as described in Example 26 to give the 7-hydroxy derivative (0.13g, 15%). Alkylation of this compound with 80% sodium hydride 0.25 mmol and benzyl bromide (0.22 mmol) in THF afforded the title compound (40 mg, 34%). MH⁺ 541.

35 **Example 32-44 were prepared by the following generic procedure:**

To a stirred solution of 2-(2-piperidin-2-yl-ethyl)-1,2,3,4-tetrahydroisoquinoline (D4) (1 mmol) and diisopropylethylamine (1 mmol) in dichloromethane cooled by an ice

bath was added an aryl sulfonyl chloride (1 mmol). Stirring was continued allowing the solution to reach room temperature over 24 hours. The solution was washed thoroughly (10% NaOH) and brine, dried and concentrated *in vacuo*. The residue was purified by chromatography on silica gel.


5



Example	Ar	MH ⁺
32	4,5-Dibromo-2-thiophene	549
33	2-Bromophenyl	463/465
34	3-Bromophenyl	463/465
35	4-Bromophenyl	463/465
36	2-Naphthyl	435
37	4-Iodophenyl	510
38	4-tert Butylphenyl	441
39	4-n-Propylphenyl	427
40	3-Methylphenyl	399
41	4-Chloro-2,5-dimethylphenyl	447
42	4-Cyanophenyl	410
43	3-Chloro-4-methyl	433
44	3,4-Dibromophenyl	541/543/545

Examples 45-52 were prepared according to the procedure outlined in Example 32-44 using 7-phenyl-2-(2-piperidin-2-yl-ethyl)-1,2,3,4-tetrahydroisoquinoline and an aryl sulfonyl chloride.

10



Example	Ar	MH ⁺
45	2-Naphthyl	511

46	3,4-Dichlorophenyl	529/531
47	4,5-Dibromo-2-thiophene	623/625/627/629
48	4-Bromophenyl	539/541
49	3-Chloro-4-methylphenyl	509/511
50	4-Chloro-3,6-dimethylphenyl	523/525
51	4-tert Butylphenyl	517
52	3-Bromophenyl	539/541

Pharmacological Data

[³H]-5-Carboxamidotryptamine binding to human 5-HT ₇ receptor clones expressed in 293 cells *in vitro*.

The affinity of test drugs for the 5-HT ₇ receptor binding site can be determined by assessing their ability to displace [³H]-5-carboxamidotryptamine from 5-HT ₇ receptor clones expressed in 293 cells (To *et al.*, 1995 and Sleight *et al.*, 1995).

10

The cells suspension (400μl) was incubated with [³H]-5-carboxamido-tryptamine (0.5nM) in Tris HCl buffer (pH 7.4) at 37°C for 45mins. Non-specific binding was measured in the presence of 5-hydroxytryptamine (10⁻⁶M). Ten concentrations of test drug (10⁻¹¹ to 10⁻⁵M final concentration) were added in a volume of 50ul. The total assay volume was 500μl. Incubation was stopped by rapid filtration using a Tomtec cell harvester and radioactivity measured by scintillation counting on a Packard Topcount. The IC₅₀ values and pK_i values were calculated by INFLEXION, a non-linear iterative curve fitting programme based in EXCEL (Bowen and Jerman, 1994).

15


- 20 Bowen, W. and Jerman, J. (1994). Br. J. Pharmacol., 112, 440P.
 Sleight, A.J., Carolo, C., Petit, N., Zweingelstein, C. and Bourson, A. (1995). Mol. Pharmacol., 47, 99.
 To, Z.P., Bonhaus, D.W., Eglen, R.M. and Jakeman, L.B. (1995). Br. J. Pharmacol., 15, 107.

25

All the compounds of examples 1 to 52 showed activity in the above test.

Claims:

1. A compound of formula (I) or a salt thereof:



(I)

- 10 wherein:

Ar is an optionally substituted mono- or bicyclic aromatic or heteroaromatic ring;

Ar' is an optionally substituted mono- or bicyclic aromatic or heteroaromatic ring;

R¹ is C₁-6alkyl or together with the group R³ form a 5-8 membered ring containing one or two heteroatoms optionally substituted by C₁-6alkyl;

15 R² is hydrogen or C₁-6alkyl;

R³ is hydrogen, C₁-6alkyl or together with the group R¹ form a 5-8 membered ring containing one or two heteroatoms optionally substituted by C₁-6alkyl;

R⁴ is hydrogen or C₁-6 alkyl;

R⁵ and R⁶ are independently hydrogen or C₁-6alkyl;

20 p is 1, 2 or 3;

q is 1 to 3; and

r is 1 or 2.

2. A compound according to claim 1 in which Ar is an optionally substituted bicyclic aromatic group.

- 25 3. A compound according to any one of claims 1 or 2 in which Ar' is an optionally substituted monocyclic aromatic group.

4. A compound according to claim 1 which is:

2-(2-[1-Naphthalene-1-sulfonyl]piperidin-2-yl)-ethyl-1,2,3,4-tetrahydroisoquinoline

2-(2-(3-Chloro-4-methylphenyl)-piperidin-2-yl)-ethyl-1,2,3,4 tetrahydroisoquinoline

- 30 and pharmaceutically acceptable salts thereof.


5. A compound according to any one of claims 1 to 4 for use in therapy.
6. A pharmaceutical composition which comprises a compound according to any one of claims 1 to 4 and a pharmaceutically acceptable carrier or excipient.
7. A process for the preparation of a compound of formula (I) or a pharmaceutically acceptable salt thereof, which process comprises:
- 5 (a) the coupling of a compound of formula (II):



10

(II)


in which Ar is as defined in formula (I) and L is a leaving group with a compound of formula (III):



15


(III)

- in which p, q, r, R², R³, R⁴, R⁵, R⁶ and Ar' are as defined in formula (I);
- 20 or (b) the coupling of a compound of formula (IV):



(IV)

- 25 in which Ar, R¹, R², R³, R⁵ and R⁶ are as defined in formula (I) and L¹ is a leaving group with a compound of formula (V):



- 5 in which q , r , R^4 and Ar' are as defined in formula (I) and optionally thereafter (a) or (b):
- forming a pharmaceutically acceptable salt.
- 8 Use of a compound according to any one of claims 1 to 4 for the manufacture
- 10 of a medicament for the treatment of anxiety and/or depression.

INTERNATIONAL SEARCH REPORT

In. tional Application No
PCT/EP 97/03160

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D401/06 C07D401/14 C07D409/14 A61K31/47

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 306 375 A (SYNTHELABO) 8 March 1989 see page 19; claims ---	1,5,6,8
A	EP 0 076 072 A (BEECHAM WUELFING GMBH & CO KG) 6 April 1983 cited in the application see claims ---	1,5,6
A	EP 0 064 445 A (CHOAY SA) 10 November 1982 see claims -----	1,5,6,8

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- * & * document member of the same patent family

Date of the actual completion of the international search

12 September 1997

Date of mailing of the international search report

19.09.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Henry, J

INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP 97/03160

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.: 1-3, 5-8
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
The wordings "optionnaly substituted mono or bicyclic aromatic or hetero-aromatic ring" are too broadly formulated to permit an adequate search.
The search has therefore been restricted to these compounds of formula I which are supported by the examples.
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

In International Application No

PCT/EP 97/03160

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0306375 A	08-03-89	FR 2619110 A	10-02-89
		FR 2625504 A	07-07-89
		FR 2630113 A	20-10-89
		FR 2630114 A	20-10-89
		AU 598149 B	14-06-90
		AU 2047588 A	09-02-89
		FR 2634204 A	19-01-90
		FR 2634205 A	19-01-90
		JP 1100167 A	18-04-89
		US 4885302 A	05-12-89
		US 4945096 A	31-07-90
EP 0076072 A	06-04-83	AU 8862282 A	31-03-83
		CA 1173443 A	28-08-84
		JP 58065267 A	18-04-83
		US 4485108 A	27-11-84
EP 0064445 A	10-11-82	FR 2504528 A	29-10-82
		AT 9996 T	15-11-84
		CA 1196919 A	19-11-85
		JP 1053669 B	15-11-89
		JP 57181053 A	08-11-82

1000
1000


1998:42391 Document No. 128:102010 Preparation of sulfonamide derivatives as 5-HT₇ receptor antagonists. Forbes, Ian Thomson; Rahman, Shirley Katherine (Smithkline Beecham P.L.C., UK; Forbes, Ian Thomson; Rahman, Shirley Katherine). PCT Int. Appl. WO9749695 A1 971231, 20 pp. DESIGNATED STATES: W: JP, US; RW: AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE. (English). CODEN: PIXXD2. APPLICATION: 97WO-EP03160 970617. PRIORITY: 96GB-0013261 960625; 96GB-0013262 960625.

IT ***201135-27-5P*** ***201135-30-0P*** ***201135-31-1P*** ***201135-32-2
P*** ***201135-38-8P*** ***201135-39-9P*** ***201135-51-5P***
201135-53-7P ***201135-54-8P***

(prepn. of sulfonamide derivs. as 5-HT₇ receptor antagonists)

RN 201135-27-5 ZCAPLUS


CN Piperidine, 2-[2-(3,4-dihydro-7-nitro-2(1H)-isoquinoliny)ethyl]-1-(1-naphthalenylsulfonyl)- (9CI) (CA INDEX NAME)



1000-1000
1000-1000

RN 201135-30-0 ZCAPLUS


CN Piperidine, 2-[2-(3,4-dihydro-6-methoxy-2(1H)-isoquinoliny)ethyl]-1-(1-naphthalenylsulfonyl)- (9CI) (CA INDEX NAME)



RN 201135-31-1 ZCAPLUS


CN Piperidine, 2-[2-(3,4-dihydro-7-methoxy-2(1H)-isoquinoliny)ethyl]-1-(1-naphthalenylsulfonyl)- (9CI) (CA INDEX NAME)

RN 201135-31-1 ZCAPLUS




RN 201135-32-2 ZCAPLUS

CN Acetamide, *N*-[1,2,3,4-tetrahydro-2-[2-[1-(1-naphthalenylsulfonyl)-2-piperidiny]ethyl]-5-isoquinoliny]-
(9CI) (CA INDEX NAME)



RN 201135-38-8 ZCAPLUS


CN Piperidine, 2-[2-(7-amino-3,4-dihydro-2(1*H*)-isoquinoliny)ethyl]-1-(1-naphthalenylsulfonyl)- (9CI) (CA
INDEX NAME)



RN 201135-39-9 ZCAPLUS


CN Piperidine, 2-[2-(7-bromo-3,4-dihydro-2(1*H*)-isoquinoliny)ethyl]-1-(1-naphthalenylsulfonyl)- (9CI) (CA
INDEX NAME)

RN 201135-39-9 ZCAPLUS




RN 201135-51-5 ZCAPLUS

CN Piperidine, 2-[2-(3,4-dihydro-6-hydroxy-2(1*H*)-isoquinolinyl)ethyl]-1-(1-naphthalenylsulfonyl)- (9CI)
(CA INDEX NAME)



RN 201135-53-7 ZCAPLUS


CN Piperidine, 2-[2-(5-amino-3,4-dihydro-2(1*H*)-isoquinolinyl)ethyl]-1-(1-naphthalenylsulfonyl)- (9CI) (CA INDEX NAME)



RN 201135-54-8 ZCAPLUS

CN Piperidine, 2-[2-(5-bromo-3,4-dihydro-2(1*H*)-isoquinolinyl)ethyl]-1-(1-naphthalenylsulfonyl)- (9CI) (CA INDEX NAME)

RN 201135-54-8 ZCAPLUS




IT ***201135-26-4P*** ***201135-28-6P*** ***201135-29-7P*** ***201135-33-3
P*** . ***201135-34-4P*** ***201135-35-5P*** ***201135-36-6P***
201135-37-7P ***201135-40-2P*** ***201135-41-3P*** ***201135-42-4
P*** ***201135-43-5P*** ***201135-44-6P*** ***201135-45-7P***
201135-46-8P ***201135-47-9P*** ***201135-48-0P*** ***201135-49-1
P*** ***201135-50-4P*** ***201135-52-6P*** ***201135-55-9P***
201135-56-0P ***201135-57-1P*** ***201135-58-2P*** ***201135-59-3
P*** ***201135-60-6P*** ***201135-61-7P*** ***201135-62-8P***
201135-63-9P ***201135-64-0P*** ***201135-65-1P*** ***201135-66-2
P*** ***201135-67-3P*** ***201135-68-4P*** ***201135-69-5P***
201135-70-8P ***201135-71-9P*** ***201135-72-0P*** ***201135-73-1
P*** ***201135-74-2P*** ***201135-75-3P*** ***201135-76-4P***
201135-77-5P

(prepn. of sulfonamide derivs. as 5-HT₇ receptor antagonists)

RN 201135-26-4 ZCAPLUS


CN Piperidine, 2-[2-(3,4-dihydro-2(1H)-isoquinolinyl)ethyl]-1-(1-naphthalenylsulfonyl)- (9CI) (CA INDEX NAME)



RN 201135-28-6 ZCAPLUS


CN Piperidine, 2-[2-[3,4-dihydro-7-(methylsulfonyl)-2(1H)-isoquinolinyl]ethyl]-1-(1-naphthalenylsulfonyl)- (9CI) (CA INDEX NAME)

RN 201135-28-6 ZCAPLUS




RN 201135-29-7 ZCAPLUS

CN 7-Isoquinolinesulfonamide, 1,2,3,4-tetrahydro-2-[2-[1-(1-naphthalenylsulfonyl)-2-piperidyl]ethyl]- (9CI)
(CA INDEX NAME)



RN 201135-33-3 ZCAPLUS


CN Piperidine, 2-[2-(8-chloro-3,4-dihydro-2(1H)-isoquinoliny)ethyl]-1-(1-naphthalenylsulfonyl)- (9CI) (CA INDEX NAME)



RN 201135-34-4 ZCAPLUS


CN Piperidine, 2-[2-(3,4-dihydro-6,7-dimethoxy-2(1H)-isoquinoliny)ethyl]-1-(1-naphthalenylsulfonyl)- (9CI)
(CA INDEX NAME)

RN 201135-34-4 ZCAPLUS




RN 201135-35-5 ZCAPLUS

CN Piperidine, 2-[2-(7-benzoyl-3,4-dihydro-2(1H)-isoquinolinyl)ethyl]-1-(1-naphthalenylsulfonyl)- (9CI)
(CA INDEX NAME)



RN 201135-36-6 ZCAPLUS


CN Piperidine, 2-[2-(3,4-dihydro-7-phenoxy-2(1H)-isoquinolinyl)ethyl]-1-(1-naphthalenylsulfonyl)- (9CI)
(CA INDEX NAME)



RN 201135-37-7 ZCAPLUS


CN Piperidine, 2-[2-(3,4-dihydro-8-phenyl-2(1H)-isoquinolinyl)ethyl]-1-(1-naphthalenylsulfonyl)- (9CI) (CA INDEX NAME)

RN 201135-37-7 ZCAPLUS




RN 201135-40-2 ZCAPLUS

CN Piperidine, 2-[2-(3,4-dihydro-7-phenyl-2(1H)-isoquinolinyl)ethyl]-1-(1-naphthalenylsulfonyl)- (9CI) (CA INDEX NAME)



RN 201135-41-3 ZCAPLUS


CN Piperidine, 2-[2-[3,4-dihydro-7-(2-methylphenyl)-2(1H)-isoquinolinyl]ethyl]-1-(1-naphthalenylsulfonyl)- (9CI) (CA INDEX NAME)



RN 201135-42-4 ZCAPLUS

CN Piperidine, 2-[2-[3,4-dihydro-7-(4-methylphenyl)-2(1H)-isoquinolinyl]ethyl]-1-(1-naphthalenylsulfonyl)- (9CI) (CA INDEX NAME)


RN 201135-42-4 ZCAPLUS



RN 201135-43-5 ZCAPLUS

CN Piperidine,


2-[2-[3,4-dihydro-7-[4-(trifluoromethyl)phenyl]-2(1H)-isoquinolinyl]ethyl]-1-(1-naphthalenylsulfonyl)- (9CI)
(CA INDEX NAME)



RN 201135-44-6 ZCAPLUS

CN Piperidine,


2-[2-[3,4-dihydro-7-[3-(trifluoromethyl)phenyl]-2(1H)-isoquinolinyl]ethyl]-1-(1-naphthalenylsulfonyl)- (9CI)
(CA INDEX NAME)



RN 201135-45-7 ZCAPLUS


CN Piperidine, 2-[2-[3,4-dihydro-7-(3-methoxyphenyl)-2(1H)-isoquinolinyl]ethyl]-1-(1-naphthalenylsulfonyl)-
(9CI) (CA INDEX NAME)

RN 201135-45-7 ZCAPLUS




RN 201135-46-8 ZCAPLUS

CN Piperidine, 2-[2-[3,4-dihydro-7-(3-pyridinyl)-2(1H)-isoquinolinyl]ethyl]-1-(1-naphthalenylsulfonyl)- (9CI)
(CA INDEX NAME)



RN 201135-47-9 ZCAPLUS


CN Piperidine, 2-[2-[3,4-dihydro-7-(2-methoxyphenyl)-2(1H)-isoquinolinyl]ethyl]-1-(1-naphthalenylsulfonyl)- (9CI) (CA INDEX NAME)



RN 201135-48-0 ZCAPLUS


CN Piperidine, 2-[2-[3,4-dihydro-7-(4-methoxyphenyl)-2(1H)-isoquinolinyl]ethyl]-1-(1-naphthalenylsulfonyl)- (9CI) (CA INDEX NAME)

RN 201135-48-0 ZCAPLUS




RN 201135-49-1 ZCAPLUS

CN Piperidine, 2-[2-[7-(3-chlorophenyl)-3,4-dihydro-2(1H)-isoquinolinyl]ethyl]-1-(1-naphthalenylsulfonyl)-
(9CI) (CA INDEX NAME)



RN 201135-50-4 ZCAPLUS


CN Piperidine, 2-[2-[3,4-dihydro-7-(trifluoromethyl)-2(1H)-isoquinolinyl]ethyl]-1-(1-naphthalenylsulfonyl)-
(9CI) (CA INDEX NAME)



RN 201135-52-6 ZCAPLUS


CN Piperidine, 2-[2-(3,4-dihydro-6-phenyl-2(1H)-isoquinolinyl)ethyl]-1-(1-naphthalenylsulfonyl)- (9CI) (CA INDEX NAME)

RN 201135-52-6 ZCAPLUS




RN 201135-55-9 ZCAPLUS

CN Piperidine, 2-[2-(3,4-dihydro-5-phenyl-2(1H)-isoquinolinyl)ethyl]-1-(1-naphthalenylsulfonyl)- (9CI) (CA INDEX NAME)



RN 201135-56-0 ZCAPLUS


CN Piperidine, 2-[2-[3,4-dihydro-7-(phenylmethoxy)-2(1H)-isoquinolinyl]ethyl]-1-(1-naphthalenylsulfonyl)- (9CI) (CA INDEX NAME)



RN 201135-57-1 ZCAPLUS


CN Piperidine, 1-[(4,5-dibromo-2-thienyl)sulfonyl]-2-[2-(3,4-dihydro-2(1H)-isoquinolinyl)ethyl]- (9CI) (CA INDEX NAME)

RN 201135-57-1 ZCAPLUS




RN 201135-58-2 ZCAPLUS

CN Piperidine, 1-[(2-bromophenyl)sulfonyl]-2-[2-(3,4-dihydro-2(1H)-isoquinolinyl)ethyl]- (9CI) (CA INDEX NAME)



RN 201135-59-3 ZCAPLUS


CN Piperidine, 1-[(3-bromophenyl)sulfonyl]-2-[2-(3,4-dihydro-2(1H)-isoquinolinyl)ethyl]- (9CI) (CA INDEX NAME)



RN 201135-60-6 ZCAPLUS


CN Piperidine, 1-[(4-bromophenyl)sulfonyl]-2-[2-(3,4-dihydro-2(1H)-isoquinolinyl)ethyl]- (9CI) (CA INDEX NAME)

RN 201135-60-6 ZCAPLUS




RN 201135-61-7 ZCAPLUS

CN Piperidine, 2-[2-(3,4-dihydro-2(1*H*)-isoquinolinyl)ethyl]-1-(2-naphthalenylsulfonyl)- (9CI) (CA INDEX NAME)



RN 201135-62-8 ZCAPLUS


CN Piperidine, 2-[2-(3,4-dihydro-2(1*H*)-isoquinolinyl)ethyl]-1-[(4-iodophenyl)sulfonyl]- (9CI) (CA INDEX NAME)



RN 201135-63-9 ZCAPLUS


CN Piperidine, 2-[2-(3,4-dihydro-2(1*H*)-isoquinolinyl)ethyl]-1-[[4-(1,1-dimethylethyl)phenyl]sulfonyl]- (9CI) (CA INDEX NAME)

RN 201135-63-9 ZCAPLUS




RN 201135-64-0 ZCAPLUS

CN Piperidine, 2-[2-(3,4-dihydro-2(1H)-isoquinolinyl)ethyl]-1-[(4-propylphenyl)sulfonyl]- (9CI) (CA INDEX NAME)



RN 201135-65-1 ZCAPLUS


CN Piperidine, 2-[2-(3,4-dihydro-2(1H)-isoquinolinyl)ethyl]-1-[(3-methylphenyl)sulfonyl]- (9CI) (CA INDEX NAME)



RN 201135-66-2 ZCAPLUS


CN Piperidine, 1-[(4-chloro-2,5-dimethylphenyl)sulfonyl]-2-[2-(3,4-dihydro-2(1H)-isoquinolinyl)ethyl]- (9CI) (CA INDEX NAME)

RN 201135-66-2 ZCAPLUS




RN 201135-67-3 ZCAPLUS

CN Piperidine, 1-[(4-cyanophenyl)sulfonyl]-2-[2-(3,4-dihydro-2(1*H*)-isoquinoliny)ethyl]- (9CI) (CA INDEX NAME)



RN 201135-68-4 ZCAPLUS


CN Piperidine, 1-[(3-chloro-4-methylphenyl)sulfonyl]-2-[2-(3,4-dihydro-2(1*H*)-isoquinoliny)ethyl]- (9CI) (CA INDEX NAME)



RN 201135-69-5 ZCAPLUS


CN Piperidine, 1-[(3,4-dibromophenyl)sulfonyl]-2-[2-(3,4-dihydro-2(1*H*)-isoquinoliny)ethyl]- (9CI) (CA INDEX NAME)

RN 201135-69-5 ZCAPLUS




RN 201135-70-8 ZCAPLUS

CN Piperidine, 2-[2-(3,4-dihydro-7-phenyl-2(1H)-isoquinolinyl)ethyl]-1-(2-naphthalenylsulfonyl)- (9CI) (CA INDEX NAME)



RN 201135-71-9 ZCAPLUS


CN Piperidine, 1-[(3,4-dichlorophenyl)sulfonyl]-2-[2-(3,4-dihydro-7-phenyl-2(1H)-isoquinolinyl)ethyl]- (9CI) (CA INDEX NAME)



RN 201135-72-0 ZCAPLUS


CN Piperidine, 1-[(4,5-dibromo-2-thienyl)sulfonyl]-2-[2-(3,4-dihydro-7-phenyl-2(1H)-isoquinolinyl)ethyl]- (9CI) (CA INDEX NAME)

RN 201135-72-0 ZCAPLUS




RN 201135-73-1 ZCAPLUS

CN Piperidine, 1-[(4-bromophenyl)sulfonyl]-2-[2-(3,4-dihydro-7-phenyl-2(1H)-isoquinolinyl)ethyl]- (9CI)
(CA INDEX NAME)



RN 201135-74-2 ZCAPLUS


CN Piperidine, 1-[(3-chloro-4-methylphenyl)sulfonyl]-2-[2-(3,4-dihydro-7-phenyl-2(1H)-isoquinolinyl)ethyl]- (9CI) (CA INDEX NAME)



RN 201135-75-3 ZCAPLUS


CN Piperidine, 1-[(4-chloro-2,5-dimethylphenyl)sulfonyl]-2-[2-(3,4-dihydro-7-phenyl-2(1H)-isoquinolinyl)ethyl]- (9CI)
(CA INDEX NAME)

RN 201135-75-3 ZCAPLUS




RN 201135-76-4 ZCAPLUS

CN Piperidine, 2-[2-(3,4-dihydro-7-phenyl-2(1*H*)-isoquinoliny)ethyl]-1-[[4-(1,1-dimethylethyl)phenyl]sulfonyl]- (9CI) (CA INDEX NAME)



RN 201135-77-5 ZCAPLUS

CN Piperidine, 1-[(3-bromophenyl)sulfonyl]-2-[2-(3,4-dihydro-7-phenyl-2(1*H*)-isoquinoliny)ethyl]- (9CI) (CA INDEX NAME)




IT ***201135-79-7P*** ***201135-80-0P***
(prepn. of sulfonamide derivs. as 5-HT₇ receptor antagonists)

RN 201135-79-7 ZCAPLUS


CN Methanesulfonic acid, trifluoro-,
1,2,3,4-tetrahydro-2-[2-[1-(1-naphthalenylsulfonyl)-2-piperidinyl]ethyl]-6-isoquinoliny ester (9CI) (CA INDEX NAME)

RN 201135-79-7 ZCAPLUS





RN 201135-80-0 ZCAPLUS

CN Piperidine, 2-[2-(3,4-dihydro-7-hydroxy-2(1H)-isoquinoliny)ethyl]-1-(1-naphthalenylsulfonyl)- (9CI)
(CA INDEX NAME)



GI



AB The title compds. [I; Ar = (un)substituted mono- or bicyclic arom. or heteroarom. ring; Ar¹ = (un)substituted mono- or bicyclic arom. or heteroarom. ring; R¹ = C₁₋₆ alkyl; R¹R³ = (un)substituted 5-8 membered ring contg. 1-2 heteroatoms; R² = H, C₁₋₆ alkyl; R³ = H, C₁₋₆ alkyl; R⁴ = H, C₁₋₆ alkyl; R⁵, R⁶ = H, C₁₋₆ alkyl; p = 1-3; q = 1-3; r = 1-2], useful in

RN 201135-80-0 ZCAPLUS

the treatment of CNS disorders such as anxiety and/or depression, were prepd. Thus, reaction of 2-(2-chloroethyl)-1-(naphthalene-1-sulfonyl)piperidine with 1,2,3,4-tetrahydroisoquinoline in the presence of NaI and K_2CO_3 in MeCN afforded 40% the title compd. II. Compds. I are effective at 0.5-100 mg/day.